



Welcome to AP Physics 2021-2022! Your summer assignment will include some textbook readings and practice problems. We will be using an online textbook for this class, so please follow the steps below in order to sign up, login and download your text book.


1. Sign up for OpenStax: Go to the following website URL: openstax.org You should land on a page that looks like the image below. I highly recommend that you sign up using your JBHA google account.




2. Now select Subjects, then Science at the top left hand corner of the screen.



3. The textbook you will use is the second option, "College Physics: for AP Courses".



4. You can either download a copy or use the online version. I recommend downloading the textbook.



5. Your summer assignment is to **read and take notes on sections 11.1-11.5 and 11.7 (pg 446-459, 464-470)**. Make sure to check out each worked-out example given. This is where you will get all of your information for this material. Complete the following problems at the end of the chapter on pages 485-490. For density values, use the table on pg. 448. You may look up the mass of any common object such as the sun, an atom, etc.

Conceptual Questions:

1-24, every 6th question (1, 6, 12, 18, 24)

30

Problems & Exercises

4, 6, 10

11, 13

14, 16, 20

25, 26

37, 39, 44, 47

Feel free to email me if you run into any problems or have any questions at dwilson@jbha.org. I have attached the AP Physics formula sheet below. You will have access to this on every assignment next year. Have a great summer!

ADVANCED PLACEMENT PHYSICS 2 EQUATIONS, EFFECTIVE 2015

CONSTANTS AND CONVERSION FACTORS	
Proton mass, $m_p = 1.67 \times 10^{-27}$ kg	Electron charge magnitude, $e = 1.60 \times 10^{-19}$ C
Neutron mass, $m_n = 1.67 \times 10^{-27}$ kg	1 electron volt, $1 \text{ eV} = 1.60 \times 10^{-19}$ J
Electron mass, $m_e = 9.11 \times 10^{-31}$ kg	Speed of light, $c = 3.00 \times 10^8$ m/s
Avogadro's number, $N_0 = 6.02 \times 10^{23}$ mol ⁻¹	Universal gravitational constant, $G = 6.67 \times 10^{-11}$ m ³ /kg·s ²
Universal gas constant, $R = 8.31$ J/(mol·K)	Acceleration due to gravity at Earth's surface, $g = 9.8$ m/s ²
Boltzmann's constant, $k_B = 1.38 \times 10^{-23}$ J/K	
1 unified atomic mass unit,	$1 \text{ u} = 1.66 \times 10^{-27}$ kg = 931 MeV/c ²
Planck's constant,	$h = 6.63 \times 10^{-34}$ J·s = 4.14×10^{-15} eV·s
	$hc = 1.99 \times 10^{-25}$ J·m = 1.24×10^3 eV·nm
Vacuum permittivity,	$\epsilon_0 = 8.85 \times 10^{-12}$ C ² /N·m ²
Coulomb's law constant, $k = 1/4\pi\epsilon_0 = 9.0 \times 10^9$ N·m ² /C ²	
Vacuum permeability,	$\mu_0 = 4\pi \times 10^{-7}$ (T·m)/A
Magnetic constant, $k' = \mu_0/4\pi = 1 \times 10^{-7}$ (T·m)/A	
1 atmosphere pressure,	$1 \text{ atm} = 1.0 \times 10^5$ N/m ² = 1.0×10^5 Pa

UNIT SYMBOLS	meter, m	mole, mol	watt, W	farad, F
	kilogram, kg	hertz, Hz	coulomb, C	tesla, T
	second, s	newton, N	volt, V	degree Celsius, °C
	ampere, A	pascal, Pa	ohm, Ω	electron volt, eV
	kelvin, K	joule, J	henry, H	

PREFIXES		
Factor	Prefix	Symbol
10^{12}	tera	T
10^9	giga	G
10^6	mega	M
10^3	kilo	k
10^{-2}	centi	c
10^{-3}	milli	m
10^{-6}	micro	μ
10^{-9}	nano	n
10^{-12}	pico	p

VALUES OF TRIGONOMETRIC FUNCTIONS FOR COMMON ANGLES							
θ	0°	30°	37°	45°	53°	60°	90°
$\sin \theta$	0	1/2	3/5	$\sqrt{2}/2$	4/5	$\sqrt{3}/2$	1
$\cos \theta$	1	$\sqrt{3}/2$	4/5	$\sqrt{2}/2$	3/5	1/2	0
$\tan \theta$	0	$\sqrt{3}/3$	3/4	1	4/3	$\sqrt{3}$	∞

The following conventions are used in this exam.

- I. The frame of reference of any problem is assumed to be inertial unless otherwise stated.
- II. In all situations, positive work is defined as work done on a system.
- III. The direction of current is conventional current: the direction in which positive charge would drift.
- IV. Assume all batteries and meters are ideal unless otherwise stated.
- V. Assume edge effects for the electric field of a parallel plate capacitor unless otherwise stated.
- VI. For any isolated electrically charged object, the electric potential is defined as zero at infinite distance from the charged object.

ADVANCED PLACEMENT PHYSICS 2 EQUATIONS, EFFECTIVE 2015

MECHANICS

$$v_x = v_{x0} + a_x t$$

$$x = x_0 + v_{x0} t + \frac{1}{2} a_x t^2$$

$$v_x^2 = v_{x0}^2 + 2a_x(x - x_0)$$

$$\vec{a} = \frac{\sum \vec{F}}{m} = \frac{\vec{F}_{net}}{m}$$

$$|\vec{F}_f| \leq \mu |\vec{F}_n|$$

$$a_c = \frac{v^2}{r}$$

$$\vec{p} = m\vec{v}$$

$$\Delta \vec{p} = \vec{F} \Delta t$$

$$K = \frac{1}{2} mv^2$$

$$\Delta E = W = F_{\parallel} d = Fd \cos \theta$$

$$P = \frac{\Delta E}{\Delta t}$$

$$\theta = \theta_0 + \omega_0 t + \frac{1}{2} \alpha t^2$$

$$\omega = \omega_0 + \alpha t$$

$$x = A \cos(\omega t) = A \cos(2\pi f t)$$

$$x_{cm} = \frac{\sum m_i x_i}{\sum m_i}$$

$$\vec{\alpha} = \frac{\sum \vec{\tau}}{I} = \frac{\vec{\tau}_{net}}{I}$$

$$\tau = r_{\perp} F = r F \sin \theta$$

$$L = I \omega$$

$$\Delta L = \tau \Delta t$$

$$K = \frac{1}{2} I \omega^2$$

$$|\vec{F}_s| = k |\vec{x}|$$

a = acceleration
 A = amplitude
 d = distance
 E = energy
 F = force
 f = frequency
 I = rotational inertia
 K = kinetic energy
 k = spring constant
 L = angular momentum
 ℓ = length
 m = mass
 P = power
 p = momentum
 r = radius or separation
 T = period
 t = time
 U = potential energy
 v = speed
 W = work done on a system
 x = position
 y = height
 α = angular acceleration
 μ = coefficient of friction
 θ = angle
 τ = torque
 ω = angular speed

$$U_s = \frac{1}{2} k x^2$$

$$\Delta U_g = mg \Delta y$$

$$T = \frac{2\pi}{\omega} = \frac{1}{f}$$

$$T_s = 2\pi \sqrt{\frac{m}{k}}$$

$$T_p = 2\pi \sqrt{\frac{\ell}{g}}$$

$$|\vec{F}_g| = G \frac{m_1 m_2}{r^2}$$

$$\vec{g} = \frac{\vec{F}_g}{m}$$

$$U_G = -\frac{G m_1 m_2}{r}$$

ELECTRICITY AND MAGNETISM

$$|\vec{F}_E| = \frac{1}{4\pi\epsilon_0} \frac{|q_1 q_2|}{r^2}$$

$$\vec{E} = \frac{\vec{F}_E}{q}$$

$$|\vec{E}| = \frac{1}{4\pi\epsilon_0} \frac{|q|}{r^2}$$

$$\Delta U_E = q \Delta V$$

$$V = \frac{1}{4\pi\epsilon_0} \frac{q}{r}$$

$$|\vec{E}| = \left| \frac{\Delta V}{\Delta r} \right|$$

$$\Delta V = \frac{Q}{C}$$

$$C = \kappa \epsilon_0 \frac{A}{d}$$

$$E = \frac{Q}{\epsilon_0 A}$$

$$U_C = \frac{1}{2} Q \Delta V = \frac{1}{2} C (\Delta V)^2$$

$$I = \frac{\Delta Q}{\Delta t}$$

$$R = \frac{\rho \ell}{A}$$

$$P = I \Delta V$$

$$I = \frac{\Delta V}{R}$$

$$R_s = \sum_i R_i$$

$$\frac{1}{R_p} = \sum_i \frac{1}{R_i}$$

$$C_p = \sum_i C_i$$

$$\frac{1}{C_s} = \sum_i \frac{1}{C_i}$$

$$B = \frac{\mu_0}{2\pi} \frac{I}{r}$$

A = area
 B = magnetic field
 C = capacitance
 d = distance
 E = electric field
 \mathcal{E} = emf
 F = force
 I = current
 ℓ = length
 P = power
 Q = charge
 q = point charge
 R = resistance
 r = separation
 t = time
 U = potential (stored) energy
 V = electric potential
 v = speed
 κ = dielectric constant
 ρ = resistivity
 θ = angle
 Φ = flux

$$\vec{F}_M = q\vec{v} \times \vec{B}$$

$$|\vec{F}_M| = |q\vec{v}| |\sin \theta| |\vec{B}|$$

$$\vec{F}_M = I\vec{\ell} \times \vec{B}$$

$$|\vec{F}_M| = |I\vec{\ell}| |\sin \theta| |\vec{B}|$$

$$\Phi_B = \vec{B} \cdot \vec{A}$$

$$\Phi_B = |\vec{B}| \cos \theta |\vec{A}|$$

$$\mathcal{E} = -\frac{\Delta \Phi_B}{\Delta t}$$

$$\mathcal{E} = B \ell v$$

ADVANCED PLACEMENT PHYSICS 2 EQUATIONS, EFFECTIVE 2015

FLUID MECHANICS AND THERMAL PHYSICS

$\rho = \frac{m}{V}$ $P = \frac{F}{A}$ $P = P_0 + \rho gh$ $F_b = \rho Vg$ $A_1 v_1 = A_2 v_2$ $P_1 + \rho gy_1 + \frac{1}{2} \rho v_1^2 = P_2 + \rho gy_2 + \frac{1}{2} \rho v_2^2$ $\frac{Q}{\Delta t} = \frac{kA \Delta T}{L}$ $PV = nRT = Nk_B T$ $K = \frac{3}{2} k_B T$ $W = -P \Delta V$ $\Delta U = Q + W$	A = area F = force h = depth k = thermal conductivity K = kinetic energy L = thickness m = mass n = number of moles N = number of molecules P = pressure Q = energy transferred to a system by heating T = temperature t = time U = internal energy V = volume v = speed W = work done on a system y = height ρ = density
--	--

MODERN PHYSICS

$E = hf$ $K_{\max} = hf - \phi$ $\lambda = \frac{h}{p}$ $E = mc^2$	E = energy f = frequency K = kinetic energy m = mass p = momentum λ = wavelength ϕ = work function
--	---

WAVES AND OPTICS

$\lambda = \frac{v}{f}$ $n = \frac{c}{v}$ $n_1 \sin \theta_1 = n_2 \sin \theta_2$ $\frac{1}{s_i} + \frac{1}{s_o} = \frac{1}{f}$ $ M = \left \frac{h_i}{h_o} \right = \left \frac{s_i}{s_o} \right $ $\Delta L = m\lambda$ $d \sin \theta = m\lambda$	d = separation f = frequency or focal length h = height L = distance M = magnification m = an integer n = index of refraction s = distance v = speed λ = wavelength θ = angle
--	--

GEOMETRY AND TRIGONOMETRY

Rectangle $A = bh$ Triangle $A = \frac{1}{2}bh$ Circle $A = \pi r^2$ $C = 2\pi r$	A = area C = circumference V = volume S = surface area b = base h = height ℓ = length w = width r = radius
--	---

Rectangular solid $V = \ell wh$ Cylinder $V = \pi r^2 \ell$ $S = 2\pi r \ell + 2\pi r^2$ Sphere $V = \frac{4}{3} \pi r^3$ $S = 4\pi r^2$	Right triangle $c^2 = a^2 + b^2$ $\sin \theta = \frac{a}{c}$ $\cos \theta = \frac{b}{c}$ $\tan \theta = \frac{a}{b}$
--	--

